

PHY138Y Mechanics - Written Homework

This homework assignment is due by 5PM on Friday October 13. It should be submitted in the *Drop Box* for your tutorial. The Drop Boxes are located in the basement of the Burton tower of McLennan. On the first floor of McLennan there is a stairway with a bust of Newton beside it; the Drop Boxes are at the bottom of that stairway.



This page must be the first page of the submitted Problem Set.



You must solve the problems together in the same team that you have been working with in your tutorials.



Problem Sets done by a single individual will not be accepted.



By filling out and signing the form below, I certify that I took an active role in the solution of all the problems of this problem set.

Name (Please Print)	Signature


Designate one member of your team as the *coordinator*. The coordinator will be responsible for assembling the final copy of your solutions and submitting them in the Drop Box on time.

The coordinator should fill out this form:

Name of Your PHY138 Tutor	
Tutorial Group	
Tutorial Day	
Tutorial Time	
Coordinator's (your) Name	

Problem 1 (15 Points)

In class on Wednesday, September 27, we analysed the situation shown to the right. Two blocks of masses M and m are stacked on top of each other and are being pulled to the right on a rough table top by a force F . The frictional force exerted on the lower block is f . The acceleration of the blocks is a . We assumed the F is greater than f , so the direction of the acceleration is as shown.



When we considered the two blocks together as the *system* we showed that the acceleration is:


$$a = \frac{F - f}{M + m}$$

We then analysed the upper block as the system. The only horizontal force being exerted on it is the static frictional one being exerted on it by the lower block. That force points to the right and "drags" the block along with the lower one. We called that force f' and showed that:

$$f' = m a$$

Here the acceleration is just the result found above.

Now you will analyse the system of the lower block alone. In a naive analysis where one is, for example, unaware of Newton's 3rd Law, one might write down a Free Body Diagram of the horizontal forces acting on the block as shown.

**Part A (7 Points)**

Show that this Free Body Diagram gives the wrong value of the acceleration

Part B (8 Points)

Now draw the correct Free Body Diagram and solve for the acceleration. Compare the result to the acceleration given above: if they are the same then your answer is correct.

Problem 2 (10 Points)

A Flash animation based on the Air Track Experiment in the labs is available via the button to the right.



If you are reading this in hardcopy, the URL is:

<http://faraday.physics.utoronto.ca/PVB/Harrison/Flash/ClassMechanics/AirTrack/AirTrack.html>

Part A (5 Points)

Verify that the total momentum is conserved for all 6 possible collisions.

Part B (5 Points)

Leibniz, a contemporary and rival of Newton, analysed collisions in terms of the *vis viva*, literally the *living force*. It is equal to the mass times the instantaneous speed squared: mv^2 . Some of the air track collisions conserve the total vis viva and some do not. Determine which.

Problem 3 (15 Points)

In class we discussed the *ballistocardiogram*. When the heart beats it expels a mass m of blood into the aorta with speed v , and the body and platform move in the opposite direction with speed V . The blood velocity can be determined independently by observing the Doppler shift of ultrasound; the Doppler effect will be studied in the next quarter of PHY138. In one trial v is 50.0 cm/s. The mass of the subject plus the platform is 54.0 kg. The platform moves 6×10^{-5} m in 0.160 s after one heartbeat.

Part A (13 Points)

Calculate the mass m of blood that leaves the heart. Assume that the mass of the blood is negligible compared to the mass of the person.

Part B (2 Points)

From your answer to Part A, is the assumption that you were asked to make reasonable? What other assumptions have you made in solving this problem?

Problem 4 (10 Points)

This question is from a recent *Medical College Admission Test* (MCAT) which many of you will be taking at some time in the future. On the MCAT it is in multiple choice format.

A projectile with a mass of 0.2 kg and a horizontal speed of 2.0 m/s hits a recycle bin (which is free to move), then rebounds at 1.0 m/s back along the same path. What is the magnitude of the horizontal momentum the bin receives?

Problem 5 (25 Points)

Knight Chapter 8 Problem 48.

Problem 6 (25 Points)

Knight Chapter 9 Problem 72. Assume, reasonably, that as the chair slides across the floor the force of friction on it is constant and equal to 39 N.